

[Sign In](#) | [My EndNote Web](#) | [My ResearcherID](#) | [My Citation Alerts](#) | [My Saved Searches](#) | [Log Out](#) | [Help](#)

ISI Web of Knowledge™

Web of Science

Additional Resources

[Search](#) | [Cited Reference Search](#) | [Advanced Search](#) | [Search History](#) | [Marked List \(0\)](#)

Web of Science®

<< [Back to results list](#)

Record 8 of 10

Record from Web of Science®

Effects of reverse osmosis isolation on reactivity of naturally occurring dissolved organic matter in physicochemical processes

[Print](#) | [E-mail](#) | [Add to Marked List](#)
[NCBI](#) | [Save to EndNote Web](#)
[Save to EndNote, RefMan, ProCite](#) [more options](#)

Author(s): Kilduff JE, Mattaraj S, Wigton A, Kitis M, Karanfil T

Source: WATER RESEARCH **Volume:** 38 **Issue:** 4
Pages: 1026-1036 **Published:** FEB 2004

Times Cited: 21 **References:** 27 [Citation Map](#)

Abstract: A field reverse osmosis system was used to isolate dissolved organic matter (DOM) from two lacustrine and two riverine surface water sources. The rejection of DOM was on the order of 99% and did not vary significantly with pressure. A simple mass balance model using a single measured value of rejection predicted the concentration within the closed-loop isolation system. The effect of operating pressure and solution flux on mass recovery of DOM was evaluated in laboratory and field trials. Under controlled laboratory conditions, >99% of a lacustrine DOM could be accounted for. A fraction of the isolated DOM was not recoverable using hydrodynamic cleaning; however, this fraction was recovered by using a pH 10 NaOH wash solution. The mass recovered in the NaOH solution increased from <1% to >6% with increasing transmembrane pressures from 414 kPa (60 psi) to 1000 kPa (145 psi), respectively. This is consistent with fouling that results from an increase in solution flux, and a decrease in tangential crossflow velocity. Under field conditions, mass balances were generally >95% and mass recovery was >90% in all cases. The effects of temperature on solution flux were consistent with changes in fluid viscosity; effects of temperature on membrane diffusivity or morphological properties were small. RO isolation under low pressure conditions designed to maximize DOM recovery had little effect on DOM reactivity evaluated in terms of nanofiltration membrane fouling, XAD-8 resin adsorption, activated carbon adsorption, competition with trichloroethylene for adsorption sites on activated carbon, and molecular weight distribution measured using size exclusion chromatography. (C) 2003 Elsevier Ltd. All rights reserved.

Document Type: Article

Language: English

Author Keywords: reverse osmosis; natural organic matter; isolation; organic carbon; fouling; nanofiltration; preloading; size exclusion chromatography

Cited by: 21

This article has been cited 21 times (from Web of Science).

Mattaraj S, Phimpha W, Hongthong P, et al. [Effect of operating conditions and solution chemistry on model parameters in crossflow reverse osmosis of natural organic matter](#) DESALINATION 253 1-3 38-45 APR 2010

Yuan YX, Kilduff JE [Hydrodynamic Modeling of NOM Transport in UF: Effects of Charge Density and Ionic Strength on Effective Size and Sieving](#) ENVIRONMENTAL SCIENCE & TECHNOLOGY 43 14 5449-5454 JUL 15 2009

An D, Song JX, Gao W, et al. [Molecular weight distribution for NOM in different drinking water treatment processes](#) DESALINATION AND WATER TREATMENT 5 1-3 267-274 MAY 2009

[[view all 21 citing articles](#)]

[Create Citation Alert](#)

Related Records:

Find similar records based on shared references (from Web of Science).

[[view related records](#)]

References: 27

View the bibliography of this record (from Web of Science).

Suggest a correction

If you would like to improve the quality of this product by suggesting corrections, [please fill](#)

KeyWords Plus: AQUATIC HUMIC SUBSTANCES;
DISINFECTION BY-PRODUCTS; ACTIVATED CARBON; TCE
ADSORPTION; FLUX DECLINE; ULTRAFILTRATION;
NANOFILTRATION; MEMBRANES; SURFACE; WATERS

out this form.

Reprint Address: Kilduff, JE (reprint author), Rensselaer
Polytech Inst, Dept Civil & Environm Engr, 110,8th St, Troy, NY
12180 USA

Addresses:

1. Rensselaer Polytech Inst, Dept Civil & Environm Engr, Troy,
NY 12180 USA
2. Ubon Ratchathani Univ, Fac Engr Chem Engr, Ubon
Ratchathani 34190, Thailand
3. Suleyman Demirel Univ, Dept Environm Engr, TR-32200
Isparta, Turkey
4. Clemson Univ, Dept Environm Sci & Engr, Anderson, SC
29625 USA

E-mail Addresses: kilduff@rpi.edu

Publisher: PERGAMON-ELSEVIER SCIENCE LTD, THE
BOULEVARD, LANGFORD LANE, KIDLINGTON, OXFORD
OX5 1GB, ENGLAND

Subject Category: Engineering, Environmental; Environmental
Sciences; Water Resources

IDS Number: 775WY

ISSN: 0043-1354

DOI: 10.1016/j.watres.2003.10.049

[<< Back to results list](#)

◀ Record 8 of 10 ▶

Record from **Web of Science®**

Output Record

Step 1:

- Authors, Title, Source
 plus Abstract
 Full Record
 plus Cited Reference

Step 2:

[\[How do I export to bibliographic management software?\]](#)

[Print](#) [E-mail](#) [Add to Marked List](#)
[Save to EndNote Web](#) [Save to EndNote, RefMan, ProCite](#)

Save to other Reference Software

[Save](#)

View in [简体中文](#) [English](#) [日本語](#)

Please give us your [feedback](#) on using ISI Web of Knowledge.

[Acceptable Use Policy](#)
 Copyright © 2010 Thomson Reuters



THOMSON REUTERS

Published by Thomson Reuters